

Unmasking Health Determinants and Health Outcomes for Urban First Nations Using Respondent Driven Sampling

Firestone M., Smylie J., Maracle S., Spiller M., O'Campo P.

Corresponding Author:

Firestone, Michelle

Post-Doctoral Fellow, Centre for Research on Inner City Health (CRICH), St. Michael's Hospital

Smylie, Janet

Research Scientist, Centre for Research on Inner City Health (CRICH)

Associate Professor, Department of Public Health Sciences, University of Toronto
Toronto, Canada

Maracle, Sylvia

Executive Director, Ontario Federation of Indian Friendship Centres (OFIFC)
Toronto, Canada

Spiller, Michael

Cornell University, Department of Sociology
Ithaca, New York

O'Campo, Patricia

Director, Centre for Research on Inner City Health

Professor, Dalla Lana School of Public Health, University of Toronto
Toronto, Canada

Keywords: Canada; urban Aboriginal health; First Nations; Respondent-Driven Sampling; Community-based Research

Word Count: 4,233

ABSTRACT

Objective: Population based health information on urban Aboriginal populations in Canada is limited due to challenges with the identification of Aboriginal persons in existing health datasets. The main objective of the Our Health Counts (OHC) project was to work in partnership with Aboriginal stakeholders to generate a culturally relevant, representative baseline health data set for three urban Aboriginal communities in Ontario, Canada. **Design:** Respondent-Driven Sampling.

Setting: Hamilton, Ontario, Canada.

Participants: The OHC study, in partnership with De dwa da dehs ney>s Aboriginal Health Access Centre (DAHC) recruited 554 First Nations adults living in Hamilton using RDS.

Results: Among First Nations adults living in Hamilton, 78% earned less than \$20,000 per year and 70% lived in the lowest income quartile neighbourhoods. Mobility and crowded living conditions were also highly prevalent. Common chronic diseases included arthritis, hypertension, diabetes and COPD and rates of emergency room access were elevated.

Conclusions: RDS is an effective sampling method in urban Aboriginal contexts as it builds upon existing social networks and successfully identified a population-based cohort. The findings illustrate striking disparities in health determinants and health outcomes between urban First Nations individuals and the general population which have important implications for health services delivery, programming and policy development.

ARTICLE SUMMARY

Strengths and limitations of this study:

- Our study is the first RDS of self-identified First Nations people residing in an urban setting in Canada
- This study models best practices in academic-Aboriginal community research partnerships
- RDS has been demonstrated to be a promising tool for generating Aboriginal health assessment measures in urban areas where there are limited existing sampling frame options
- At present, multivariable regression analyses using RDS samples have not appropriately addressed the co-relation between observations and the unequal

sampling probabilities inherent in RDS; therefore we present prevalence estimates for which methods are better established

INTRODUCTION

According to the 2006 Census, over 60% (150,570 people) of Aboriginal^a people living in Ontario live in urban areas.[1] Nearly seven out of ten Métis live in urban areas, and about three out of every four people in the off-reserve First Nations population live in urban areas.[1] First Nations people have historically been dislocated from their original homelands and may constitute diasporic, heterogeneous communities in urban areas.[2] Specifically, an increasing number of First Nations individuals are moving to urban centers to seek better housing, employment and education opportunities and for the services and amenities available.[3-5] There is a growing body of literature exploring the complexities of urban indigeneity [6] and some of the elements which distinguish urban Aboriginal identity are social and economic marginalization, a growing middle class, and population diversity in terms of cultural origins and legal status.[7] According to the Canadian Census, First Nations, Inuit, and Métis populations experience ongoing disparities in social determinants of health such as income insecurity, unemployment, low levels of education, decreased food availability, and inadequate housing compared to non-Aboriginal Canadians, and these disparities are exacerbated with urban residence.[1;8;9]

Despite the growing size of the urban Aboriginal population in Canada, accessible and culturally relevant population health data for this population are almost non-existent.[10-13] While census data do exist, there are serious deficits in population health measures.[13] Some of the reasons behind this deficiency are limitations in the current health information system and data collection processes with respect to Aboriginal peoples.[14;15] Sampling frames are often biased and comprise non-random subpopulations such as lists of members or clients of particular programs and services in the community. When urban Aboriginal people have been included in census-based national surveys, these surveys were underpowered and often First Nations, Inuit, and Métis data could not be successfully disaggregated.[14;16] Additionally, studies based on Census data show that a significant number of Aboriginal people move from rural and reserve areas to cities, and back and forth as well as within and among cities.[4;17] These factors make it difficult to get reliable counts of the Aboriginal population of a city.

^a “Aboriginal people” is a collective name for all of the original peoples of Canada and their descendants. The *Constitution Act* of 1982 specifies that the Aboriginal Peoples in Canada consist of three groups: Indians, Inuit and Métis. The term First Nations came into common use in the 1970s to replace Indian, which some people found offensive. Despite its widespread use, there is no legal definition for this term in Canada. In the United States, the term “Native American” is used to represent Indigenous peoples from North America, including Mexico. For alignment with the US Constitution and to ensure policy implications, the term “American Indian” is still used for Indigenous populations living within US borders, particularly the lower 48 states and parts of Alaska. “Indigenous” is a term used in the international context.

At the national level, the continued inability to identify Aboriginal peoples in healthcare databases leads to very poor coverage of Aboriginal populations in Canada, with a specific paucity of health information for First Nations, Métis, and Inuit living in urban areas. Provincial and territorial systems generally do not collect ethnic-specific utilization data, and their ethnic flags for vital statistics are inconsistent or absent.[18] The federal government's decision to cancel the long form of the Canadian national census will further challenge the ability of researchers and communities to identify and address social inequities across and within populations who experience racialization and the negative effects of structural discrimination in healthcare and other sectors, thus widening the existing health and social policy vacuum.[19;20]

In the absence of an accessible and accurate population based sampling frame for urban Aboriginal people, we implemented respondent driven sampling (RDS) for the Our Health Counts (OHC) study. RDS, like other chain-referral approaches, relies on members of a population to recruit their peers.[21;22] However, RDS differs from other methods because of estimation procedures that adjust for the participants' different probabilities of being sampled.[22] RDS also differs from other chain-referral methods because the number of recruitments any participant can make is limited, and it employs a dual incentive structure where participants are rewarded both for participating and for recruiting their peers.[23]

There is limited literature on RDS in Indigenous contexts;[24] however, there is evidence that links the success of RDS to the strength of social networks and relationships.[25] Indigenous knowledges are wholistic, relational and embraces a fluidity that allows for constant growth and change.[26;27] Aboriginal identities tend to value the group over the individual, thereby establishing models of kinship whereby everyone has the right to give and receive according their own choices.[26] Therefore, we hypothesized that RDS would be an appropriate and effective sampling methodology as it builds on social networks and would draw on existing kinship systems known to be present in Indigenous communities.[28]

The main objective of the OHC project was to work in partnership with Aboriginal stakeholders to generate a culturally relevant, representative baseline health data set for three urban Aboriginal communities in Ontario, Canada. The results presented here pertain specifically to the urban First Nations population in Hamilton where the organizational project lead was the Ontario Federation of Indian Friendship Centres (OFIFC) and the community partner was De dwa da dehs ney>s Aboriginal Health Access Centre (DAHC).

METHODS

Community-based participatory research

This study upheld current ethical and scientific standards in Indigenous health research[29-33] through a community based participatory research partnership between the academic research team, OFIFC and DAHC. We ensured that our Aboriginal decision making partners were active in all aspects of the research through the following mechanisms: Aboriginal leadership; research agreements which explicitly addressed

issues of project governance, community expectations, benefits, ownership, control, access, and possession of information, and dissemination of project results; capacity building through staffing at community sites, data workshops and awareness building; respect for the individual and collective rights of Aboriginal peoples with respect to their health information; cultural relevance through the development and application of culturally appropriate measures; representation of the urban Aboriginal population of Ontario; and sustainability of the project to ensure expansion of the database. In addition, a Governing Council, comprised of representatives from the core partner urban Aboriginal provincial organizations was established to oversee all stages of the research process. Specifically, this body adhered to governance protocols and ensured that individual and collective community rights were respected, were kept informed about the project's progress and led the project towards meaningful results, acted as a resource to the community on questions related to various portions of the research project and controlled the release of all data generated by the study.

All members of the study team were from the Aboriginal community, living in the general vicinity of the city of Hamilton. Formal ethics approval was provided by the Research Ethics Board of St. Michael's Hospital in Toronto.

Study Site

The City of Hamilton is located in southern Ontario on what was traditionally Haudenosaunee (Iroquian) territory and is near two First Nations reserves: Six Nations of the Grand River and Mississaugas of the New Credit. According to the 2006 Census, the total Aboriginal population in Hamilton was 13,735 people, comprising 2.8% of the total population of the city.[34] The OFIFC identified the city of Hamilton as a promising community site based on its significant Aboriginal population and strong infrastructure of Aboriginal community services.

Recruitment

The RDS process was initiated through the purposive selection of individuals to begin recruitment, also called "seeds." Inclusion criteria for participation in the study included 18 years of age, residence within the geographic boundaries of the City of Hamilton and self-identified as having First Nations identity. First Nations identity was determined through open dialogue around family, experiences, and location while also explaining the overall study goals and additional study sites in Ottawa. Questions such as: "What words best describe you?" and "How do you see yourself?" were useful. With the help of community partners, six seeds were identified and agreed to participate. In order for seeds to reflect a diverse demographic of First Nations people living in Hamilton, factors such as gender, age, family size, and occupations were considered in their selection. For example, seeds were identified in the student population, among the steel workers union, among artists, elders, housing and social assistance providers as well as members of local Aboriginal organizations and boards. Five of the six seeds produced referrals within the two weeks leading up to the December 2009 holiday closure at DAHC. In February and March 2010, three additional seeds were added.

Study participants, including seeds, provided informed consent and then completed a health assessment survey after which they received 3-5 coded coupons to refer a member of their social network into the study. Following the holiday closure of DAHC, the number of distributed coupons per recruit was increased from 3 to 5 in order to increase the rate of recruitment. Participants received a monetary reimbursement of \$20 for their time and participation. They also received an additional \$10 for every eligible person they recruited into the study. Study recruitment and interviews took place in an Aboriginal community centre.

For sample size calculation, we assumed 95% confidence intervals (CI), survey item prevalence ranges from 10% to 75% and a design effect of 2.[35] Based on this formula, the OCH study aimed to recruit 500 First Nations adults and 250 children.

Sources of Data

Community concept mapping was implemented to develop a community specific survey for First Nations in Hamilton, [36] which was comprised of 8 domains covering areas such as sociodemographics, physical, mental, and emotional health, First Nations identity and impacts of colonization (i.e. questions about Residential School attendance, involvement of child protection agencies, dislocation from traditional lands and discrimination), and access and barriers to care. The survey tool was piloted with First Nations community members, and minor modifications were made to improve face validity. Data were collected and compiled by computer-assisted personal interviewing with a computer based version of the survey tool developed by the research team using SPSS Data Collection Author and Collector.[37]

Upon completion of recruitment and survey, our sample of First Nations adults was linked to data holdings at the Institute of Clinical Evaluative Sciences (ICES) including provincial records of emergency room and hospital visits (Canadian Institutes of Health Information – Discharge Abstract Database and National Ambulatory Care Reporting System: Emergency), and neighbourhood income quintiles (derived from Census data). Using health care system identification numbers (i.e. Ontario Health Card) provided by participants, we successfully linked 92% (N=725) of First Nations adults to the ICES database.

Analysis

Recruitment dynamics, adjusted population based estimates and confidence intervals were calculated using the RDS-I enhanced data smoothing estimator in the custom RDSAT software (version 7.1).[38] The RDS methodology anticipates that personal networks are not randomly distributed, and therefore adjusts for small to moderate levels of network clustering (people who have ties to others like them), in the form of post-sampling weights.

RESULTS

Among First Nations adults in Hamilton, 78.9% of participants were recruited via referral trees originating from two seeds. With 19 and 32 waves respectively, the lengths of both of these recruitment chains were long enough to overcome the original sampling bias, which usually happens after 6 or 7 waves of recruitment.[39] The statistical theory is that if the chain-referral process consists of enough waves or cycles of recruitment, the composition of the final sample with respect to key characteristics and behaviors will become independent of the seeds from which it began. The point at which the sample composition becomes stable is termed “equilibrium” and is an indication that RDS has been implemented successfully.[21]

Over a period of four and a half months (November 2009 to March 2010) a total of 790 persons were recruited, including 554 adults and 236 children Overall, a high response rate for survey questions was observed. Specifically, non-response rates ranged from 2.5% to less than 0.5%, including sensitive questions around income, food security, and impacts of colonization, which had non-response rates under 2%. Such high response rates can be attributed to the survey tool itself, which reflected the health priorities of the community[40] and which was administered in a safe and culturally secure context. Demographic and social characteristics are presented in Table 1.

Table 1. Demographic and social characteristics of First Nations adults (Total N=554)

Characteristic	Prevalence % (95% CI)
Sex	
Male	59.9 (53.3, 67.2)
Age	
18-34	40.2 (33.3, 48)
35-49	35.4 (28.7, 41.6)
50+	24.4 (18.4, 30.8)
Education	
Some high school or less	57.3 (51.1, 64)
Completed high school	19.5 (14.5, 24.7)
Some or completed college	18.3 (13.3, 23.3)
Some or completed University	5 (2.4, 8)
Income sources for household	
Provincial or municipal social assistance or welfare (e.g. ODSP, Ontario Works)	69.2 (63.1, 75)
Wages and salaries	28.2 (22.1, 34.2)
Child Tax Benefit	17 (12.4, 21.8)
Any other income support	13.3 (9.2, 17.6)
Employment insurance	9.6 (5.4, 14.3)
Income from self-employment	7.7 (4.3, 11.7)
Child Support payments	3 (1.1, 4.6)
Individual annual income	
\$0-4,999	18.3 (13.4, 24.8)
\$5,000-5,999	23.1 (17.1, 28.5)

\$10,000-14,999	21.9 (16.3, 27.6)
\$15,000-19,999	14.8 (10.7, 20)
\$20,000+	21.8 (16, 27.6)
Mobility	
No moves in past 5 years	10.2 (6.2, 14.8)
1 move in past 5 years	15.5 (10.5, 20.6)
2 moves in past 5 years	20.2 (15.4, 26.4)
3-5 moves in past 5 years	41.1 (35, 48.6)
6-10 moves in past 5 years	10.4 (5.7, 13.4)
Overcrowding*	
<=1 persons per room	27.7 (21.6, 33.8)
1-2 persons per room	46 (40.5, 54.3)
>2 persons per room	26.3 (19.7, 30.8)
Food security	
You and others always had enough of the kinds of food you wanted to eat	26.7 (21.1, 32.8)
You and others had enough to eat, but not always the kinds of food you wanted	51.5 (45.3, 58)
Sometimes or often you did not have enough to eat**	21.8 (16.5, 27)

Note. CI = confidence interval.

*Following Statistics Canada standards: calculated by dividing the number of rooms in each household (excluding the bathroom) by the number of people residing in the home, where a higher value of 'persons per room' indicates a higher level of crowding[41]

** Collapsed sometimes and often you did not have enough food to eat

The data presented in Table 2 for age, gender and income quintiles compare the OHC population estimates with the total Hamilton population and a random subset of 10% of the Ontario provincial population. The three populations were very similar with respect to the gender breakdown, however, the OHC sample was younger than the general Hamilton and Ontario populations which is consistent with Census data.[42] Over 70% of the First Nations population fell into the lowest income quartile compared to 25% of the general Hamilton population and 20% of the Ontario population. At the higher end of the income scale, we observed that while 15% of Hamilton residents and 20% of the Ontario population fell into the highest income quartile, only 3% of the study population was earning in this category.

Table 2. Gender, Age and Income Quintile for First Nations Adults, City of Hamilton and Ontario-10%

Variable	Sample			
	OHC (Total N=725*)	OHC-RDS- Adjusted	Hamilton (Total N=536,253)	Ontario-10% (Total N=1,324,241)
	Prevalence, %	Prevalence, % (95% CI)	Prevalence, %	Prevalence, %

Sex				
Female	49.4	37.6 (29.6, 43.6)	50.9	51.2
Male	50.6	62.4 (56.4, 70.4)	49.1	48.7
Age**				
18-34	37.4	41.9 (34.4, 49.9)	28.2	28.2
35-49	37.6	36.6 (29.9, 43.1)	28.1	29.5
50-64	22.9	20.7 (14.7, 26.9)	24.8	24.8
65+	2.1	0.8 (0.3, 1.6)	18.9	17.6
Income Quintile				
1-Low	71.76	73 (66.5, 79.2)	25.1	19.5
2	16.22	11.8 (7.7, 16)	22.8	19.8
3	6.3	7.4 (3.6, 10.5)	20.3	19.8
4	2.48	4.9 (2.5, 9.8)	17	20.4
5-High	1.34	3 (1.1, 5.4)	14.7	20.2

Note. CI = confidence interval; OHC = Our Health Counts; RDS = Respondent Driven Sampling.

*Total N reflects number of participants linked to ICES database

**Age of participants on 2010-04-01

The most common self-reported chronic conditions that had been diagnosed by a health care provider among First Nations adults (n=554) in Hamilton included arthritis (30.7%; 95% CI 25.4-36.8), hypertension (25.8%; 95% CI 20.3-31.6), asthma (19%; 95% CI 14.9-24.6), diabetes (15.6%; 95% CI 11.2-21.1), heart disease (8.4%; 95% CI 4.8-12.5) and COPD (8.4%; 95% CI 5.3-11.8). The prevalence of self-reported Hepatitis C was 8.7% (95% CI 4.3-11.7). 25% reported having been injured over the past 12 months. Self-reported upper respiratory tract infection (URTI) was common, with 73% (95% CI 67.9-79.9) of adults reporting URTI in the past 12 months.

Among First Nations adults (n=554), 16.7% (95% CI 11.4 – 22.1) felt that availability of health services in their community was excellent, 43.3% (95% CI 36.1 - 49.6) felt it was good, 28.9% (95% CI 23.3 – 35.2) felt it was fair and 11.1% (95% CI 7.9 – 15.6) felt availability was poor. The fact that 40% of the population felt their level of access to health care was fair or poor, despite the geographic proximity to extensive health and social services that the City of Hamilton provides, substantiates the idea that just because the services are geographically proximate, does not mean that they are accessible to First Nations people. Close to half (47.9%) of the First Nations living in Hamilton reported that long waiting lists to see a specialist were a barrier (95% CI 41.9 - 55.4). Other common barriers included: not being able to arrange transportation; doctor not available; not being able to afford direct costs and/or transportation; services not covered by Non-Insured Health Benefits and lack of trust in health care provider.

Both acute and non-acute emergency room visits were much more frequent among First Nations in Hamilton compared to the general Hamilton and Ontario populations. According to data linkages with ICES, a striking 10.6% of the First Nations adult population in Hamilton made 6 or more emergency room visits in the previous 2

years compared to 1.6% and 1.9% of the Hamilton and Ontario adult populations respectively (Table 3).

Table 3. Number of Emergency Room visits in the Previous 2 Years for Adults (18-64 years), First Nations, City of Hamilton and Ontario-10%

Frequency of ER visits	Sample			
	OHC (Total N=514*)	OHC RDS-Adjusted	Hamilton (Total N=360,378)	Ontario-10% (Total N=901,509)
	Prevalence, %	Prevalence, % (95% CI)	Prevalence, %	Prevalence, %
None	31.5	(25.8 – 37.5)	66.3	69.1
1	26.2	(20.7 – 32.8)	18.7	16.8
2-5	31.7	(25.9 – 37.8)	13.4	12.3
6+	10.6	(6.2 – 14.5)	1.6	1.9
ER visits (acute)				
None	50.2	(43.9 – 57.5)	78.4	80.1
1	24.7	(18.7 – 30.1)	14.2	13.1
2-5	20.7	(15.3 – 26.1)	6.8	6.3
6+	4	1.6 – 6.9)	0.6	0.6
ER visits (non-acute)				
None	54.3	(47.6 – 61.2)	79.4	81.2
1	22.4	(17.3 – 28.5)	14	12.2
2-5	20.4	(14.5 – 25.4)	6.3	6
6+	2.9	(1 – 5.6)	0.4	0.6

Note. CI= confidence interval; OHC = Our Health Counts; RDS = Respondent Driven Sampling

*Total N reflects number of adults aged 28-64 linked to ICES database

Based on ICES data linkages, rates of hospitalization were slightly higher among the First Nations population compared to the Hamilton and Ontario populations. This may partly be explained by a higher birth rate among First Nations populations compared to the general Canadian population,[43] which is reflected by higher rates of obstetrics hospitalization compared to the general Hamilton and Ontario population: 7.3% (95% CI, 3.9 – 10.9) of the Hamilton First Nations population between the ages of 18 and 64 years had been hospitalized at least once in the past 5 years for obstetrical reasons compared to 4% of both the general Hamilton and Ontario populations in the same age group and over the same time period. However, given the high rates of chronic diseases such as diabetes and heart disease and the much higher rates of emergency room use among the First Nations population, these findings were unexpected and therefore require further investigation.

DISCUSSION

The OHC project generated a representative, urban Aboriginal health database. Through the successful application of RDS, a sample of urban First Nations individuals who might otherwise not have participated in mainstream surveys or the census was captured. In addition, given the current deficiency in public health information on urban First Nations populations, the statistically rigorous RDS sample allowed us to produce population-based estimates that can be utilized by local, regional, provincial and federal policy makers as well as urban Aboriginal stakeholders to address social inequities and health disparities facing this community.

Striking levels of poverty were identified in this study. The income data presented here may be a more representative income profile of the First Nations population in Hamilton than the 2006 Census. Specifically, the OHC data were adjusted for the sampling procedure using an RDS estimator and captured more ‘hidden’ populations such as persons who are homeless, transient or who have low literacy skills, who are known to be under-represented in the Census.[11;13] Further evidence of low incomes among this population is provided by the RDS-adjusted ICES income quintile data, which revealed that over 70% of the OHC cohort was in the lowest income quartile compared to 25% of the general Hamilton population and 20% of the Ontario population.

Accompanying high levels of poverty, housing and food insecurity were also highly prevalent among First Nations persons living Hamilton. Over half of the study population had moved at least 3 times in the past 5 years compared to 60% of the general Hamilton population who had not moved at all for the past 5 years.[34] Compared to an overcrowding rate of 3% for the general Canadian population in the 2006 Census,[1] 74% of First Nations persons in Hamilton live in crowded conditions. These findings echo those presented in the 2007 Urban Aboriginal Task Force (UATF) final report, which documented persistent obstacles with finding affordable housing, obtaining stable and secure employment and accessing appropriate services and resources among urban Aboriginal people.[44] Similarly, the Toronto Aboriginal Research Project (TARP) found that Aboriginal people in Toronto tend to concentrate in lower-income neighbourhoods where a significant proportion, particularly men, live in poverty.[45] Similar experiences have been reported in other urban jurisdictions across Canada.[46;47] While a growing proportion of urban Aboriginal people in Canada are achieving varying degrees of economic success in a number of occupations,[5;45] there remains a significant amount of unmet basic needs among this population.

The burden of chronic conditions experienced by First Nations adults in Hamilton is disproportionately high. For example, 19% of the OHC Hamilton population reported having been told by a health care provider that they have asthma, which is more than twice the self-reported asthma rate (9.3%) for the overall Hamilton population in the 2007 Canadian Community Health Survey (CCHS).[48] Likewise, rates of arthritis and high blood pressure were 30.7% and 25.8% among First Nations adults in Hamilton compared to 19.9% and 19.7% among all adults in Hamilton as reported in the 2007 CCHS.[48] For First Nations adults in Hamilton, the rate of diabetes diagnosed by a health care provider was approximately three times the rate among the general Hamilton population.[48] These findings are particularly disconcerting given the much younger age demographic of the First Nations Hamilton population and the documented barriers

to receiving health care services including long wait lists, challenges with accessing and affording transportation and the unavailability of primary care providers including physicians, nurses, and other healthcare providers in the area. Furthermore, 45.8% (95% CI 38.9-52.4) of First Nations adults believed that their ability to engage in preventative health activities (i.e. regular exercise, going to the doctor or nurse for health screening tests, accessing preventative dental care) had been affected by financial hardship.

High rates of emergency usage by First Nations persons living in Hamilton are likely linked to the problems in accessing non-emergency health care, as revealed by 40% of the population who rated their access to health care as fair or poor and 48% who indicated that wait lists are too long. In addition, as a result of primary healthcare reforms,[49] patients may be told by their primary care providers to go to the emergency department rather than a walk-in clinic when their primary care team is not available.[50] High rates of emergency room visits may also reflect a perceived lack of access to community-based or primary care settings and the extent to which high proportions of First Nations people continue to be dismissed or disregarded when attempting to access healthcare, despite geographic proximity. [51-53] Further examination is required to explore potential bias in hospital admission practices and to differentiate between types of hospitalizations (i.e. mental health, surgical and medical hospitalizations).

There were a number of limitations of the study. Firstly, it should be noted that we may have under sampled the elderly First Nations population living in nursing homes or residences for senior citizens in Hamilton. During our initial seed selection, we did not select any seeds from this population, and this population may have fewer network ties than other population groups, which may have resulted in an under representation of First Nations individuals over 65 years of age. Secondly, RDS cannot guarantee a random sample of network members;[54] therefore, estimates are unbiased to the degree that the assumptions of the RDS estimator are met. There is an active literature on RDS and estimation using RDS data,[55;56] and there is debate regarding the true design effects of RDS surveys.[57;58] Existing multivariable regression analyses using RDS samples have not appropriately addressed the correlation between observations and the unequal sampling probabilities inherent in RDS, therefore, we have focused on reporting prevalence estimates for which methods are better established. Despite these limitations, RDS was used to effectively recruit an urban Aboriginal population and allowed for the derivation of rigorous population estimates and data linkages that have previously not been possible.

Given the near absence of population health information for urban Aboriginal people in Canada, this research is able to provide, for the first time, First Nations data that clearly demonstrate alarming socio-economic inequities, a significant burden of chronic disease, multiple barriers in access to health care and elevated emergency room use. This newly established health database represents a significant contribution to public health that will directly inform strategic directions for the improvement of health and social status of urban Aboriginal people in Ontario. For example, this research will support the development of RDS as a tool to effectively recruit a larger cohort for longitudinal research with Aboriginal families in Ontario.

AKCNOWLEDGEMENTS

Firstly, we would like to thank the First Nations men and women who participated in the Our Health Counts project and so generously shared their time and stories over the course of several months. The Our Health Counts project would not have been possible without the tremendous commitment and enthusiasm of the research staff at De dwa da dehs ney>s Aboriginal Health Access Centre in Hamilton. This project was funded by the Ontario Federation of Indian Friendship Centres (OFIFC), the Ministry of Health and Long Term Care (MOHLTC) Aboriginal Health Transition Fund, and the Centre for Research on Inner City Health (CRICH) at St. Michael's Hospital. The Institute for Clinical Evaluative Sciences (ICES) contributed the costs of the in-house ICES data analysis. Dr. Smylie was supported by a Canadian Institutes for Health Research New Investigator in Knowledge Translation award during the course of the project.

Contributorship statement: MF contributed substantially to the study conception and design, co-led data interpretation and analysis and drafted multiple versions of the manuscript. JS co-led the study conception and design, data analysis and interpretation and made important contributions to the draft and final manuscript. SM contributed substantially to study conception and design, was a member of the governing council that guided each stage of research, represented the Aboriginal organizational lead of the research and made important revisions to the draft manuscript. MS provided important technical support during data collection, co-led the data analysis and interpretation, and provided important revisions to the manuscript. PO contributed substantially to the study conception and design, analysis and interpretation of data and made important revisions to the draft manuscript.

Competing interests: There are no competing interests.

Funding: This project was funded by the Ontario Federation of Indian Friendship Centres (OFIFC), the Ministry of Health and Long Term Care (MOHLTC) Aboriginal Health Transition Fund, and the Centre for Research on Inner City Health (CRICH) at St. Michael's Hospital.

Data sharing: No additional data are available.

Reference List

- [1] Statistics Canada. Aboriginal Peoples in Canada in 2006: Inuit, Métis and First Nations, 2006 Census. Ottawa ON: Ministry of Industry; 2008.
- [2] Canadian Institute for Health Research (CIHR). CIHR Guidelines for Health Research Involving Aboriginal People. Ottawa: CIHR; 20097.
- [3] National Aboriginal Health Organization (NAHO). Ways of Knowing: A Framework for Health Research. National Aboriginal Health Organization; 2003.
- [4] Policy Research Initiative. Not Strangers in these Parts. Urban Aboriginal Peoples. 2003.
- [5] Environics Institute. Urban Aboriginal Peoples Study Main Report. Environics Institute; 2010.
- [6] Peters E., Andersen C. Indigenous in the City: Contemporary Identities and Cultural Innovation. Vancouver: UBC Press, 2013.
- [7] Andersen C. Urban Aboriginality as a Distinctive Identity, in Twelve Parts. In: Peters E., Andersen C., editors. Indigenous in the City: Contemporary Identities and Cultural Innovation. Vancouver: UBC Press, 2013. 46.
- [8] Garner R., Carrière G., Sanmartin C., Longitudinal Health and Administrative Data Research Team. **The Health of First Nations Living Off-Reserve, Inuit, and Métis Adults in Canada: The Impact of Socio-economic Status on Inequalities in Health**. Ottawa: Statistics Canada, Health Analysis Division; 2010. Report No.: 82-622-X.
- [9] Tait H. Aboriginal Peoples Survey, 2006: Inuit Health and Social Conditions. Ottawa, ON: Statistics Canada; 2008.
- [10] Smylie J. Aboriginal infant mortality rate in Canada. *Lancet* 2012;380(9851):1384.
- [11] Anderson M., Smylie J. Performance measurement systems in Canada: How well do they perform in First Nations, Inuit and Métis Contexts? *Pimatisiwin: A Journal of Aboriginal and Indigenous Community Health* 2009;7(1):99-115.
- [12] Smylie J., Fell D., Ohlsson A. A Review of Aboriginal Infant Mortality Rates in Canada: Striking and Persistent Aboriginal and Non-Aboriginal Inequities. *Canadian Journal of Public Health* 2010;101(2):143-148.

- [13] Smylie J., Anderson M. Understanding the health of Indigenous peoples in Canada: key methodologic and conceptual challenges. *Canadian Medical Association Journal* 2006;165(6):602-605.
- [14] Smylie J., Anderson I., Ratima M., et al. Who is Measuring and Why? Indigenous Health Performance Measurement Systems in Canada, Australia, and New Zealand. *The Lancet* 2006;367:2029-2031.
- [15] Siggner A.J. The Challenge of Measuring the Demographic and Socio-Economic Conditions of the Urban Aboriginal Population. In: Newhouse D., Peters E., editors. *Strangers in these Parts: Urban Aboriginal Peoples*. Ottawa: Policy Research Initiative, 2003. 119-130.
- [16] Smylie J., Lofters A., Firestone M., O'Campo P. Population-Based Data and Community Empowerment. In: O'Campo P., Dunn J., editors. *Rethinking Social Epidemiology: Towards a Science of Change*. Springer Publishing Company, 2011.
- [17] Graham K.A., Peters E.J. *Aboriginal Communities and Urban Sustainability*. Ottawa: Canadian Policy Research Networks; 2002.
- [18] Anderson M., Smylie J., Anderson I., Sinclair R., Crengle S., Ratima M. First Nations, Métis, and Inuit Health Indicators in Canada. A background paper for the project "Action oriented indicators of health and health systems development for Indigenous peoples in Canada, Australia, and New Zealand. Saskatoon: Indigenous Peoples Health Research Centre; 2006.
- [19] Collier R. Long-form census changes worries health researchers. *Canadian Medical Association Journal* 2010;182(12):E563-E564.
- [20] Webster P.C. Aboriginal health programming under siege, critics charge. *Canadian Medical Association Journal* 2012;184(14):E739-E740.
- [21] Heckathorn D. Respondent driven sampling: a new approach to the study of hidden populations. *Social Problems* 1997;44:174-199.
- [22] Heckathorn D. Respondent-driven sampling II: Deriving Valid Population Estimates from Chain referral Samples of Hidden Populations. *Social Problems* 2002;49(1):11-34.
- [23] Heckathorn D, Seeman S, Broadhead R, et al. Extensions of Respondent Driven Sampling: A new approach to the Study of Injection Drug Users aged 18-25. *AIDS and Behavior* 2002;6(1):55-67.
- [24] Evans-Campbell T., Lindhorts T., Huang B., et al. Interpersonal Violence in the Lives of Urban American Indian and Alaska Native Women: Implications for Health, Mental Health, and Help-Seeking. *American Journal of Public Health* 2006;96(8):1416-1422.

- [25] Abdul-Quader A., Heckathorn D, Saidel T. Implementation and Analysis of Respondent Driven Sampling: Lessons from the Field. *Journal of Urban Health* 2006;83(7):1-5.
- [26] Greenwood M. Children as citizens of First Nations: Linking Indigenous health to early childhood development. *Pediatrics and Child Health* 2005;10(9):553-555.
- [27] Ermine W. Aboriginal Epistemology. In: Battiste M, Barman J., editors. *First Nations Education in Canada: The Circle Unfolds*. Vancouver: UBC Press, 1995. 101-112.
- [28] Smylie J., Kaplan-Myrth N., McShane K.E., et al. Indigenous Knowledge Translation: Baseline Findings in a Qualitative Study of the Pathways of Health Knowledge in Three Indigenous Communities in Canada. *Health Promotion Practice* 2009;10(3):436-446.
- [29] Canadian Institute of Health Research, Natural Sciences and Engineering Research Council of Canada, Social Sciences and Humanities Research Council of Canada. *Tri-Council Policy Statement: Ethical Conduct for Research Involving Humans*. 2010.
- [30] Canadian Institute for Health Research (CIHR). *CIHR Guidelines for Health Research Involving Aboriginal People*. Ottawa: CIHR; 2007.
- [31] Ball J., Janyst P. Enacting Research Ethics Partnerships with Indigenous Communities and Canada: "Do it in a Good Way.". *Journal of Empirical Research on Human Research Ethics* 2008;3(2):33-51.
- [32] First Nations Centre. *First Nations Conceptual Frameworks and Applied Models on Ethics, Privacy, and Consent in Health Research and Information*. Ottawa: National Aboriginal Health Organization (NAHO); 2006.
- [33] Ontario Federation of Indian Friendship Centres (OFIFC). *USAI Utility Self-Voicing Access Inter-relationality Reserach Framework*. 2012.
- [34] Statistics Canada. *Hamilton, Ontario (Code3525005) (table)*. 2006 Community Profiles. 2006 Census. Ottawa ON: Statistics Canada; 2007. Report No.: Catalogue no. 92-591-XWE.
- [35] Salganik M.J. Variance estimation, design effects, and sample size calculations for respondent-driven sampling. *Journal of Urban Health* 2006;83(6 (Suppl)):98-112.
- [36] Firestone M. *Our Health Counts: Unmasking Health and Social Disparities among Urban Aboriginal People in Ontario* University of Toronto; 2013.
- [37] IBM® SPSS® Data Collection Author [computer program]. Version 6.0.1. Chicago, IL: SPSS Inc.; 2000.

- [38] Respondent-Driven Sampling Analysis Tool (RDSAT) Version 7.1 [computer program]. Ithaca, New York: Cornell University; 2007.
- [39] Wejnert C. An Empirical Test of Respondent-Driven Sampling: Point Estimates, Variance, Degree Measures, and Out-of-Equilibrium Data. *Sociological Methodology* 2009;39(1):73-116.
- [40] Firestone M., Smylie J, Maracle S., et al. Concept Mapping: Application of a Community-Based Methodology in three Urban Aboriginal Populations. *American Indian Culture and Research Journal*, Under Review 2014;38(4).
- [41] Statistics Canada. Persons per room of private household. <http://www.statcan.gc.ca/concepts/definitions/dwelling-logement-07-eng.html> 2013 Available from: URL: <http://www.statcan.gc.ca/concepts/definitions/dwelling-logement-07-eng.html>
- [42] Statistics Canada no.92-594-XWE. Hamilton, Ontario (Code 3525005) (table). Aboriginal Population Profile. 2006 Census. <http://www12.statcan.ca/census-recensement/2006/dp-pd/prof/92-594/index.cfm?Lang=E> (accessed January 20, 2014) 2007 [cited 11 A.D. Apr 30];
- [43] Health Canada. A Statistical Profile on the Health of First Nations in Canada, for the Year 2000. Ottawa; 2005. Report No.: H35-4/30-2000E-PDF.
- [44] Ontario Federation of Indian Friendship Centres (OFIFC), Ontario Métis Aboriginal Association, Ontario Native Women's Association (ONWA). Urban Aboriginal Task Force Final Report. Toronto ON; 2007.
- [45] McCaskill D., FitzMaurice K., Cidro J. Toronto Aboriginal Research Project (TARP) Final Report. 2011.
- [46] Anderson A.B. Home in the City. University of Toronto Press, 2013.
- [47] Peters E. Indigeneity and marginalization: planning for and with urban Aboriginal communities in Canada. *Progress in Planning* 2005;63(4):327-404.
- [48] Statistics Canada. Canadian Community Health Survey. Ottawa ON: Statistics Canada; 2007.
- [49] Muldoon L., Rowan M.S., Geneau R., et al. Models of primary care Service delivery in Ontario: Why such diversity? *Healthcare Management Forum* 2006;19(4):18-23.
- [50] Howard M., Goertzen J., Kaczorowski J., et al. Emergency Department and Walk-in Clinic Use in Models of Primary Care Practice with Different After Hours Accessibility in Ontario. *Health Care Policy* 2008;4(1):73-88.

- [51] Browne A., Smye V., Rodney P., et al. Access to primary care from the perspective of Aboriginal patients at an urban emergency department. *Qualitative Health Research* 2011;21(3):333-348.
- [52] Browne A, Varcoe C., Wong S.T., et al. Closing the health equity gap: Evidence-based strategies for primary health care organizations. *International Journal for Equity in Health* 2012;11(59):1-15.
- [53] Kurtz D.L.M., Nyberg J.C., Van Den Tillaart S., et al. Silencing of voice: An act of structural violence: Urban Aboriginal women speak out about their experiences with health care. *Journal of Aboriginal Health* 2008;4(1):53-63.
- [54] Heimer R. Critical Issues and Further Questions About Respondent-Driven Sampling: Comment on Ramirez-Valles, *et al.* (2005). *AIDS and Behavior* 2005;9(4):403-408.
- [55] Gile K.J., Handcock M.S. Respondent-Driven Sampling: An Assessment of Current Methodology. *Sociological Methodology* 2010;40(1):285-327.
- [56] Heckathorn D. Extensions of respondent-driven sampling: analyzing continuous variables and controlling for differential recruitment. *Sociological Methodology* 2007;37(1):151-208.
- [57] Goel S., Salganik M.J. Assessing respondent-driven sampling. *Proceedings of the National Academy of Sciences of the United States of America* 2010;107(15):6743-6747.
- [58] Wejnert C., Pham H., Krishna N., et al. Estimating design effect and calculating sample size for respondent-driven sampling studies of injection drug users in the United States. *AIDS Behavior* 2012;16(4):797-806.